

UCSF

UC San Francisco Previously Published Works

Title

Neuroprotective Properties of Xenon.

Permalink

<https://escholarship.org/uc/item/6h44s12d>

Journal

Molecular neurobiology, 57(1)

ISSN

0893-7648

Authors

Maze, Mervyn
Laitio, Timo

Publication Date

2020

DOI

10.1007/s12035-019-01761-z

Peer reviewed

Dear Author

Here are the proofs of your article.

- You can submit your corrections **online** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- Please return your proof together with the permission to publish confirmation.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the journal title, article number, and your name when sending your response via e-mail, fax or regular mail.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.

Please note

Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI.

Further changes are, therefore, not possible.

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s12035-019-01761-z>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

The **printed version** will follow in a forthcoming issue.

Metadata of the article that will be visualized in OnlineFirst

1	Article Title	Neuroprotective Properties of Xenon	
2	Article Sub- Title		
3	Article Copyright - Year	Springer Science+Business Media, LLC, part of Springer Nature 2019 (This will be the copyright line in the final PDF)	
4	Journal Name	Molecular Neurobiology	
5	Corresponding Author	Family Name	Maze
6		Particle	
7		Given Name	Mervyn
8		Suffix	
9		Organization	UCSF
10		Division	Center for Cerebrovascular Research, Department of Anesthesia and Perioperative Care
11		Address	San Francisco, CA, USA
12		e-mail	Mervyn.Maze@ucsf.edu
13	Author	Family Name	Laitio
14		Particle	
15		Given Name	Timo
16		Suffix	
17		Organization	University of Turku
18		Division	Division of Perioperative Services, Intensive Care Medicine and Pain Management, Turku University Hospital
19		Address	Turku, Finland
20		e-mail	
21	Schedule	Received	29 August 2019
22		Revised	
23		Accepted	29 August 2019
24	Abstract	Note: This data is mandatory. Please provide.	
25	Keywords separated by ' - '	Xenon - Neuroprotection - Medical application	
26	Foot note information	Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.	

Neuroprotective Properties of Xenon

Mervyn Maze¹ · Timo Laitio²

Received: 29 August 2019 / Accepted: 29 August 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

Note: This data is mandatory. Please provide.

Keywords Xenon · Neuroprotection · Medical application

Introduction

Xenon is a colorless, odorless, tasteless, mono-atomic, and inert gas with a relative molecular weight of 131.3, and the empirical formula is Xe. Xenon is an extremely rare gas that represents no more than 0.0875 ppm in the atmosphere; this feature led the discoverers William Ramsay and Morris Travers to name it xenon from the Greek word “ξένος” (xenos) for stranger or foreign. Because of xenon’s rarity, it is extremely expensive to produce from the residue left from air separation units that are used to produce oxygen; therefore, its commercial applications have been limited to high-priced applications such as the ultimate “clean gas” in the electronics/semi-conductor industry, an ion propellant for space travel, and a bright lighting source, and for medical applications, notably anesthesia, imaging, and neuroprotection following acute ongoing injury.

In this review, the authors trace the development of xenon for medical applications from the physico-chemical properties to the initial preclinical studies, and conclude in randomized clinical trials (RCTs).

Inert but Biologically Active

Because xenon is enshrouded by five filled electron shells, it is incapable of covalent bonding and forming adducts under

biological conditions as electrons cannot be donated or accepted. However, because of xenon’s relatively high polarizability [1], with a value of 4 compared with 0.2 for helium, it can form dipoles and has an affinity for amino acid residues surrounding preformed hydrophobic cavities thereby changing the functional properties of neighboring proteins by London dispersion forces. In this manner, xenon has been shown to have activity on many proteins, governed mostly by the size and shape of xenon atoms.

Because of its chemical non-reactivity, xenon is not biotransformed, which results in two important features governing its future clinical use. Most xenobiotic drugs are converted into metabolites that may be toxic; as xenon is not metabolized, the dangers posed by toxic metabolites are obviated. Furthermore, xenon in the inspired and expired gases are identical, permitting recirculation of exhaled xenon and thereby limiting the need for a fresh supply of this scarce resource.

Xenon for Anesthesia

Xenon was reported to have anesthetic properties in 1951 [2] and comes closest to exhibiting all of the ideal properties of an inhaled general anesthetic (Table 1). Compared with nitrous oxide, the other non-potent gaseous anesthetic, xenon is 1.5 times more potent; xenon is more suitable than nitrous oxide for anesthesia because of its lower blood/gas solubility and the consequent extremely rapid inflow and washout from the body. Despite this, xenon is infrequently used as an anesthetic even though European market authorization has been in effect for more than a decade; low utilization is attributed to the high cost involved in manufacturing this rare gas from the atmosphere. Therefore, the expense of using xenon as an anesthetic for routine adult surgery appears not to be justified given the available alternatives [3].

✉ Mervyn Maze
Mervyn.Maze@ucsf.edu

¹ Center for Cerebrovascular Research, Department of Anesthesia and Perioperative Care, UCSF, San Francisco, CA, USA
² Division of Perioperative Services, Intensive Care Medicine and Pain Management, Turku University Hospital, University of Turku, Turku, Finland

t1.1	Table 1 Properties of an	
t1.2	“ideal” inhalation agent	Obtainable in pure form at a reasonable cost
t1.3		Inherently stable
t1.4		Not biotransformable
t1.5		Lacks organ-specific toxicity
t1.6		Minimal cardiorespiratory effects
t1.7		Non-flammable
t1.8		Low blood-gas partition coefficient for rapid uptake, elimination, and titratability
t1.9		Sufficiently potent allowing an enriched inspired oxygen concentration
t1.10		Lacks long-term adverse effects with chronic exposure
t1.11		Lacks unpleasant smell and irritation to airway
t1.12		Possesses analgesic and hypnotic properties
t1.13		Readily reversible central nervous system effects with no stimulation

68 Drugs produce the anesthetic state via interaction with re-
 69 ceptor targets which potentiate inhibitory neurotransmission
 70 and/or inhibit excitatory neurotransmission [4]. Xenon is
 71 thought to exert anesthetic action by potent non-competitive
 72 inhibition of the excitatory NMDA receptors [5] through an
 73 action at the binding site of the co-agonist, glycine [6]. Xenon
 74 also exerts potent effects on the neuronal background potassi-
 75 um channels including two-pore domain potassium channels
 76 such as TREK and TASK, which modulate neuronal excitabil-
 77 ity [7], and on ATP-sensitive potassium channels [8].

78 The first reported clinical experience with xenon for anes-
 79 thesia was published in 1951 by Cullen and Gross, who re-
 80 ported on two patients who underwent surgical procedures
 81 (orchietomy in an 81-year-old man and fallopian tube ligation
 82 in a 38-year-old woman who was 24-h postpartum) while
 83 receiving a xenon–oxygen (80:20) mixture to achieve the first
 84 plane of the third stage of anesthesia [2]. Induction was com-
 85 pleted within 5 min and both patients maintained normal
 86 blood pressure, pulse rate, and pulse character and had good
 87 color throughout the procedures. Within 2 min of
 88 discontinuing xenon, both patients were oriented to time,
 89 place, and person; several hours later, they were able to recol-
 90 lect information given to them at this time. Since then, numer-
 91 ous clinical studies have investigated the effects of xenon in
 92 humans.

93 Safety and tolerability information regarding xenon for in-
 94 halation stem mostly from published literature describing clin-
 95 ical studies investigating the anesthetic properties of xenon
 96 gas, and the publications contain only summary information.
 97 In aggregate, the literature suggests that administration of xe-
 98 non, as a general anesthetic, to patients both with and without
 99 cardiovascular disease is associated with hemodynamic

stability that is unparalleled in critical care settings. Side ef-
 fects identified in the literature that are frequently associated
 with the use of xenon gas for inhalation as a general anesthetic
 include raised intracranial pressure [9], bradycardia [3], and
 nausea and vomiting [10]. Although bradycardia is a safety
 concern identified for xenon anesthesia, if heart rate slows to
 the point that systemic blood pressure decreases, then standard
 positive chronotropic agents such as anti-muscarinic agents
 (e.g., glycopyrrolate or atropine) and β_1 adrenergic agonists
 (e.g., isoproterenol) can be administered to reverse it. Xenon is
 not known to interfere with oxygenation, but in an oxygen and
 xenon mixture, the greater the percentage of inhaled xenon
 administered to a subject, the lower the fraction of inspired
 oxygen that can be administered. Under circumstances in
 which lung oxygenation is compromised (e.g., from pulmo-
 nary edema), a higher fraction of inspired oxygen may be
 required to prevent arterial hypoxemia.

Xenon has a favorable pharmacokinetic (PK) profile for
 anesthesia with fast induction and emergence, which is inde-
 pendent of the duration of exposure. This PK effect is attrib-
 utable to its low blood-gas partition coefficient of 0.115 [11],
 which is significantly lower than those of other inhalational
 anesthetics (nitrous oxide, 0.47; sevoflurane, 0.65; desflurane,
 0.42). As xenon is excreted by the lungs with no biotransfor-
 mation by the renal or hepatic systems, it may prove to be the
 anesthetic of choice in certain circumstances when liver or
 kidney function decrements.

Xenon has an oil/water solubility coefficient of 20, which is
 the highest coefficient of all noble gases, and it is the only
 noble gas with anesthetic properties at atmospheric pressures.
 The physico-chemical properties of xenon are detailed in
 Table 2.

Cardiovascular Effects of Xenon in Patients Without Cardiac Diseases

In 1990, Lachmann and associates published a randomized
 double-blind trial comparing the efficacy and potency of xe-
 non with those of nitrous oxide, with special focus on the
 cardiovascular and respiratory systems [12]. The authors

Table 2 Physico-chemical properties: Ostwald solubility coefficients of xenon at 37 °C

Ostwald solubility coefficients (mL gas/mL liquid) at 37 °C		t2.1
Water/gas	0.075	t2.2
Oil/gas	1.8	t2.3
Blood/gas	0.115	t2.4
Oil/water	20	t2.5
Muscle/liver/kidney	0.10	t2.6
Adipose tissue	1.3	t2.7
Brain, gray substance	0.13	t2.8
Brain, white substance	0.23	t2.9

concluded that xenon is a more potent anesthetic than nitrous oxide in suppressing response to surgical stimuli and maintaining hemodynamic stability. Lachmann's group also compared the effect of xenon and nitrous oxide on the neurohumoral response and hemodynamics of 32 ASA class I–II patients, with the same protocol as described above [13]. The investigators concluded that xenon has more favorable hemodynamic, neurohumoral, and antinociceptive properties than nitrous oxide. Luttropp and associates investigated the effects of xenon on in vivo cardiac function using transesophageal echocardiography and hemodynamic measurements [14]. The fractional area in a short-axis view of the left ventricle at the level of the papillary muscles remained unchanged, suggesting that xenon anesthesia had no adverse effect on myocardial function as well as hemodynamics. The first multicenter randomized control trial, involving 224 patients in six centers, compared xenon/oxygen with isoflurane/nitrous oxide anesthesia and concluded that xenon anesthesia is as safe and effective as the isoflurane/nitrous oxide regimen, with the advantage that xenon exhibited more rapid recovery [15]. Also, significantly fewer xenon-anesthetized patients required inotropic support than the isoflurane group. In a single center study involving 160 patients, the hemodynamic effects were compared between those randomized to receive either xenon or propofol [16]. While systolic blood pressure was well maintained after induction with xenon at near baseline levels, propofol caused a significant post-induction decline in pressure that persisted throughout maintenance of general anesthesia. Heart rates were significantly lower in the patients who received xenon. In a multicenter study involving 252 patients scheduled for elective non-cardiovascular surgery, hemodynamic stability, including transesophageal echocardiography, was compared between patients randomized to receive either xenon or isoflurane [17]. While isoflurane decreased the myocardial contractile index, no such change was noted in the xenon-anesthetized patients, leading the authors to opine that xenon enables cardiovascular stability.

Cardiovascular Effects of Xenon in Patients With Cardiovascular Diseases

Ten patients after coronary artery bypass graft surgery (performed on cardiopulmonary bypass) were randomized to receive either propofol or xenon for sedation while being ventilated in the ICU [18]. The patients were crossed over to the alternative sedative after some hours. Compared with propofol sedation, xenon sedation did not change the heart rate or blood pressure; left ventricular stroke work index was similar.

Effects of xenon on hemodynamics in patients scheduled for coronary artery bypass graft surgery have also been assessed [19]. Statistically significant differences were found between the two groups' mean arterial pressure (MAP), fractional area change of the left ventricle, and end-diastolic area

of the left ventricle. Xenon decreased the MAP and fractional area change significantly less and increased end-diastolic area significantly more than nitrous oxide. In a safety and feasibility study involving 20 patients undergoing coronary artery bypass surgery, xenon, at varying concentrations (0%, 20%, 35%, and 50% v/v), was administered while on cardiopulmonary bypass [20]. Despite theoretical concerns about expansion of gas bubbles, the cerebral embolic load, measured by middle cerebral artery Doppler, was no higher in patients who received xenon. Troponin levels tended to be lower 24 h after surgery in patients who received xenon. Twenty-six patients scheduled for implantation of an internal cardioverter-defibrillator were randomized to receive either xenon or propofol (both with remifentanyl) for maintenance of general anesthesia [21]. Most of these patients had heart failure from ischemic heart disease or dilated cardiomyopathy. In contrast to propofol, surgical patients maintained on xenon had no changes in either the MAP or the left ventricular ejection fraction.

Central Nervous System Effects of Xenon

Volunteers ($n = 12$) were randomized to receive general anesthesia with xenon or propofol and the cerebral metabolic rate was assessed with the positron emission tomography (PET) ligand ^{18}F -fluorodeoxyglucose [22]. The xenon-exposed volunteers had cerebral metabolic rates globally reduced by 26% compared with those exposed to propofol alone. In another study, using ^{15}O -labeled water, the regional cerebral blood flow was monitored by PET scanning during xenon anesthesia in nine volunteers [23]. Xenon statistically significantly decreased the regional cerebral blood flow in several of the gray matter areas studied while regional cerebral blood flow increased by 22.1% ($\pm 13.6\%$) in the white matter. A follow-up PET study, involving five healthy subjects, assessed regional cerebral blood flow and regional cerebral glucose metabolism using ^{15}O -labeled water and ^{18}F -labeled fluorodeoxyglucose, respectively [24]. In general, the regional reduction in cerebral metabolism was greater than the regional decrement in cerebral blood flow. Luttrop et al. [14] investigated the effects of inhalation of 65% xenon on cerebral blood flow velocities, using Doppler sonography in 17 ASA class I patients undergoing abdominal surgery; they found that cerebral blood flow velocity was unchanged during the first 5 min of xenon anesthesia, but was significantly increased in the left and right, middle, and the right anterior cerebral arteries after 15 and 30 min. In addition, Giller et al. [25] noted that administration of 25%, 30%, or 35% of xenon for 5 min to normal volunteers resulted in an increase in cerebral blood flow, measured by Doppler velocity, in 85% of subjects and a decrease in cerebral blood flow in 15% of subjects. These findings are in contrast to the findings of the PET studies described in the preceding paragraph. Reasons for these discrepancies could

be differences in the patient populations (i.e., healthy volunteers versus patients undergoing surgery), differences in the duration of xenon administration, and differences in the methodology used to assess blood flow. In a trial involving supplementation of therapeutic hypothermia with administration of xenon to neonates suffering from hypoxic ischemic encephalopathy, Azzopardi and colleagues reported a significant reduction in seizure activity in patients randomized to receive 30% xenon [26].

249 Neuroprotection

250 Xenon is thought to exert neuroprotective action by acting as an
 251 antagonist at NMDA receptors. Excessive entry of calcium,
 252 mediated by NMDA receptors, triggers biochemical cascades
 253 that ultimately lead to neuronal cell death. NMDA-induced neu-
 254 rotoxicity is through “excitotoxicity” from overactivation of
 255 NMDA receptors that underlies the acute neuronal injury ob-
 256 served following insults such as stroke, cardiac arrest, and trau-
 257 matic brain injury. NMDA receptor antagonists are neuropro-
 258 tective in in vitro and in vivo brain injury models [27].
 Q6 259 Following the discovery that xenon inhibits NMDA receptors
 260 [5], it was shown that xenon could protect neuronal cell cultures
 261 against injury induced by NMDA, glutamate, or oxygen-
 262 glucose deprivation [28]. The same study showed xenon to be
 263 neuroprotective in vivo against neuronal injury caused by sub-
 264 cutaneous injection of *N*-methyl (D, L)-aspartate in rats.
 265 Subsequently, this finding was corroborated by Petzelt et al.,
 266 in an in vitro model of hypoxia [29] and in an in vivo model
 267 of stroke [30]. Other NMDA receptor antagonists such as ni-
 268 trous oxide, ketamine, and dizocilpine (MK-801) have intrinsic
 269 neurotoxicity, but xenon not only appears to be devoid of these
 270 neurotoxic effects but also ameliorates the injury produced by
 271 other NMDA antagonists [31]. Furthermore, xenon upregulates
 272 the transcription factor hypoxia inducible factor 1 alpha (HIF
 273 1 α) and its downstream cytoprotective effectors including eryth-
 274 ropoietin [32, 33]. Xenon has now been shown to afford neuro-
 275 protection in a variety of mammalian in vitro and in vivo models
 276 and meets the Stroke Treatment Academic Industry Roundtable
 277 recommendation for proceeding to clinical trials [34].

278 Phase II Clinical Trial in Out-of-Hospital Cardiac Arrest 279 Patients

280 Based upon successful animal studies investigating the effects
 281 of xenon in the setting of cardiac arrest [35, 36] and because of
 282 the synergistic interaction between xenon and therapeutic hy-
 283 pothemia [37, 38], the Xe-Hypotheca trial (NCT 00879892;
 284 May 2009–September 2014) was initiated at a single academ-
 285 ic site (University of Turku Hospital, Finland) to determine the
 286 feasibility and cardiac safety of inhaled xenon when added to
 287 therapeutic hypothermia for successfully resuscitated out-of-

hospital cardiac arrest (OHCA) patients [39]. Feasibility was
 established after the first 36 patients were randomized to re-
 ceive either therapeutic hypothermia alone ($n = 18$) or thera-
 peutic hypothermia in combination with xenon by inhalation
 ($n = 18$), with a target concentration of at least 40% xenon for
 24 h. In the xenon group, the median end-tidal xenon concen-
 tration was 47% and duration of xenon inhalation was 25.5 h.
 Xenon did not induce significant conduction, repolarization,
 or rhythm abnormalities. Median dose of norepinephrine dur-
 ing hypothermia was 2.95 mg in xenon-treated patients and
 5.30 mg in patients treated with therapeutic hypothermia alone
 ($p = 0.06$). Heart rate was statistically significantly lower in
 xenon-treated patients than that in patients treated with thera-
 peutic hypothermia alone ($p = 0.04$). From the initial results of
 this trial, the investigators concluded that xenon treatment in
 combination with hypothermia is feasible and has favorable
 cardiac features in OHCA patients. The Xe-Hypotheca trial
 was extended to a second site in 2013 (University of Helsinki
 Hospital, Finland) with an expanded cohort; the effect of xe-
 non on ischemic white matter damage was assessed by frac-
 tional anisotropy from diffusion tensor magnetic resonance
 imaging (MRI) [40]. Neurological outcome and mortality at
 6 months were also assessed. A total of 224 patients were
 screened for eligibility. One hundred and ten OHCA patients,
 aged 24–76 years, were randomized to receive either hypo-
 thermia treatment alone for 24 h (control group, $n = 55$) or
 inhaled xenon, administered to achieve an end-tidal xenon
 concentration of at least 40%, combined with hypothermia
 (33 °C) for 24 h (xenon group, $n = 55$). The primary endpoint
 was severity of ischemic white matter brain injury as evaluat-
 ed by fractional anisotropy from diffusion tensor MRI; MRIs
 were scheduled within 16 h after rewarming of a patient (rang-
 ing between 36 and 52 h after OHCA). Secondary endpoints
 were neurological outcome, assessed with cerebral perfor-
 mance category score (from 1 = conscious, alert, able to work,
 might have mild cognitive deficit, to 5 = death) and modified
 Rankin Scale (score from 0 = no symptoms at all to 6 = death),
 mortality at 6 months, and complication rate within 7 days of
 post-CA. However, the trial was not powered to detect statis-
 tically significant differences in clinical efficacy (i.e., mortal-
 ity at 6 months and neurological outcome) between groups.
 The primary endpoint was assessed in the complete case pop-
 ulation. Survival at 6 months and complication rate were an-
 alyzed in the intention-to-treat population. Kaplan–Meier sur-
 vival curves and a Cox proportional hazards model were used
 to compare mortality at 6 months between groups.

Of the randomized patients, six patients in the control
 group and seven patients in the xenon group were missing
 MRI data and were excluded from the complete case popula-
 tion. The mean (\pm SD) global fractional anisotropy value of all
 voxels in the xenon group (0.433 [\pm 0.028]) was significantly
 different than that in the control group (0.419 [\pm 0.033]) ($p =$
 0.03). The age-, gender-, and site-adjusted mean global

fractional anisotropy values were 3.8% higher in the xenon group than those in the control group (adjusted mean difference 0.016 [95% CI, 0.005 to 0.027]; $p = 0.006$). The severity of observed widespread injury was demonstrated; on average, 41.7% of the white matter tracts, including major commissural, associative, and projection fibers, were significantly more severely injured in the control group than in the xenon group. These fibers are involved in multiple important cognitive functions such as attention, memory, language, emotions, auditory, visual and executive processing, and motor functions of the body.

At 6 months, 75 patients (68.2%) were alive and able to provide follow-up data. In ordinal analysis of modified Rankin Scale, median (interquartile range) value was 1 (0 to 6) in the xenon group and 1 (0 to 6) in the control group (median difference = 0 [95% CI, 0 to 0]; $p = 0.68$). The Kaplan–Meier survival estimate (panel A) after 6 months was 27.7% (15/55 patients) in the xenon group and 34.5% (19/55 patients) in the control group (adjusted hazard ratio = 0.49 [95% CI, 0.23 to 1.01]; $p = 0.053$) (Fig. 1).

It was concluded that among comatose survivors of OHCA, treatment with inhaled xenon combined with hypothermia resulted in less white matter damage, as measured by fractional anisotropy of diffusion tensor MRI, than treatment with hypothermia alone. In contrast, there was no statistically

significant difference between groups in neurological outcomes or mortality at 6 months. However, the study was underpowered to detect a statistically significant difference in clinical outcome due to the rarity of severe neurological impairment in long-term survivors after CA; about 90% of CA patients who are alive at the 6-month follow-up have experienced a good neurological outcome (cerebral performance category 1–2). While there was no statistically significant difference in neurological outcomes or mortality at 6 months, unpublished data demonstrates that there was a trend to a survival benefit.

A predefined secondary objective was to assess the effect of inhaled xenon on myocardial ischemic damage [41]. Troponin-T levels were measured at hospital admission, and at 24 h, 48 h, and 72 h post-CA. Among comatose OHCA patients, inhaled xenon combined with hypothermia resulted in less severe myocardial injury than with hypothermia alone, as demonstrated by the significantly reduced release of troponin-T.

Rates of serious adverse events (SAEs) in the xenon group were not significantly different from the rates of SAEs in the standard of care group [40]. SAEs seen in both the xenon and standard of care groups include status epilepticus, acute kidney injury (in the “risk,” “injury,” or “failure,” RIFLE categories), pulmonary edema, ventricular fibrillation, ventricular tachycardia, atrial fibrillation, coronary stent thrombosis,




Fig. 1 Whole-brain fiber tractography of fractional anisotropy. Fractional anisotropy (FA) is a scalar value representing directionality of water diffusion. White matter damage leads to a loss of microstructural organization that can be quantified by the loss of directionality in the diffusion of water molecules in the white matter tracts. Using data from a diffusion tensor imaging sequence of an MRI scan performed within 72 h of rewarming, panels a–f represent sequential ascending horizontal planes of the major tracts in the brain. The visualization presents the

results of the voxel-wise tract-based spatial statistics analysis of FA values between the xenon group and the control group. Voxels with significantly ($p < 0.05$, family-wise error corrected for multiple comparisons) higher fractional anisotropy values in the xenon group were identified and are shown in red in the statistical visualization (i.e., 41.7% of all 119,013 analyzed voxels), whereas areas in which there were no significant difference in fractional anisotropy values between the groups are shown in green (modified from reference [40])

391 sepsis, pneumonia, multi-organ failure, adult respiratory dis-
 392 stress syndrome, and subarachnoid hemorrhage SAEs only ob-
 393 served in the xenon group included bradycardia treated with
 394 pacemaker ($n = 1$ event) and serious bleeding (gastrointestinal,
 395 $n = 1$ event). SAEs only observed in the standard of care
 396 group included third-degree atrioventricular block ($n = 1$
 397 event), carotid dissection ($n = 1$ event), carotid thrombosis
 398 ($n = 1$ event), and serious bleeding (intracranial, $n = 1$ event).

399 Conclusion and Potential Future Applications

400 Xenon exhibits many features of a putative neuroprotective
 401 agent with an ideal pharmacokinetic profile for use following
 402 acute neurological injury. Studies involving several different
 403 acute neurological injury models in a variety of animal species
 404 from four laboratories have consistently demonstrated the
 405 neuroprotective efficacy of xenon even when administered
 406 as long as 6 h following neurological injury. The mechanisms
 407 for neuroprotection appear to involve (i) antagonism of the
 408 NMDA receptor whose activation is pivotal for the excitotoxic
 409 damage that follows neurologic injury, and (ii) upregulation of
 410 HIF 1α and the resulting cytoprotection from erythropoietin, a
 411 downstream effector of the transcription factor.

412 A phase 2 RCT (Xe-Hypotheca) demonstrated significantly
 413 less white matter brain damage, as reflected by higher global
 414 fractional anisotropy values, in subjects randomized to re-
 415 ceive xenon by inhalation during the 24-h period of targeted
 416 temperature management.

417 The stage is now set for a pivotal phase 3 RCT,
 418 XePOHCAS (NCT03176186), to determine the efficacy
 419 (using endpoints of good functional outcome and survival at
 420 30 and 90 days), safety, and cost-effectiveness of xenon, at a
 421 dose of 50% of 1 atmosphere, in the management of post-
 422 cardiac arrest syndrome patients. The trial is likely to report
 423 its outcome in 2021 or before depending on the interim anal-
 424 ysis at the halfway point of the 1436-patient trial to be con-
 425 ducted in 7 countries in Europe and North America.

426 In the event that xenon exhibits neuroprotective efficacy in
 427 the XePOHCAS trial, then subsequent trials are likely to be
 428 conducted in other acute neurological injury settings including
 429 stroke and traumatic brain injury. As with the XePOHCAS trial,
 430 the maximal dose of xenon is likely to be restricted to $\leq 50\%$ of
 431 1 atmosphere as these patients may also have lung injury that
 432 will require an F_iO_2 of no less than 0.5 to avoid hypoxemia.

433 A further clinical application may be in pediatric surgical
 434 settings to obviate the occurrence of anesthetic-induced devel-
 435 opmental neurotoxicity (AIDN). For AIDN, xenon may be
 436 particularly effective both by reducing exposure to high con-
 437 centrations of neurotoxic volatile anesthetics by virtue of xe-
 438 non's anesthetic properties and because the neurotoxicity may
 439 be obviated by xenon's neuroprotective properties.

References

1. Dmochowski I (2009) Xenon out of its shell. *Nat Chem* 1(3):250
2. Cullen SC, Gross EG (1951) The anesthetic properties of xenon in animals and human beings, with additional observations on krypton. *Science* 113(2942):580–582
3. Law LS, Lo EA, Gan TJ (2016) Xenon anesthesia: a systematic review and meta-analysis of randomized controlled trials. *Anesth Analg* 122(3):678–697
4. Franks NP (2008) General anaesthesia: from molecular targets to neuronal pathways of sleep and arousal. *Nat Rev Neurosci* 9(5):370–386
5. Franks NP, Dickinson R, de Sousa SL, Hall AC, Lieb WR (1998) How does xenon produce anaesthesia? *Nature* 396(6709):324
6. Dickinson R, Peterson BK, Banks P, Similliss C, Martin JC, Valenzuela CA, Maze M, Franks NP (2007) Competitive inhibition at the glycine site of the N-methyl-D-aspartate receptor by the anesthetics xenon and isoflurane: evidence from molecular modeling and electrophysiology. *Anesthesiology* 107(5):756–767
7. Gruss M, Mathie A, Lieb WR, Franks NP (2004) The two-pore-domain $K(+)$ channels TREK-1 and TASK-3 are differentially modulated by copper and zinc. *Mol Pharmacol* 66(3):530–537
8. Bantel C, Maze M, Trapp S (2010) Noble gas xenon is a novel adenosine triphosphate-sensitive potassium channel opener. *Anesthesiology* 112(3):623–630
9. Plougmann J, Astrup J, Pedersen J, Gyldensted C (1994) Effect of stable xenon inhalation on intracranial pressure during measurement of cerebral blood flow in head injury. *J Neurosurg* 81(6):822–828
10. Lo EA, Law LS, Gan TJ (2016) Paradox of the incidence of post-operative nausea and vomiting after xenon-based anaesthesia. *Br J Anaesth* 116(6):881–883
11. Goto T, Suwa K, Uezono S, Ichinose F, Uchiyama M, Morita S (1998) The blood-gas partition coefficient of xenon may be lower than generally accepted. *Br J Anaesth* 80(2):255–256
12. PMID: 27199321 (Lo, 2016, Br J Anaesth, Paradox of the incidence of postoperative nausea and vomiting after xenon-based anaesthesia) Need PMID for Lachmann 1990
13. Boomsma F, Rupprecht J, Man in 't Veld AJ, de Jong FH, Dzoljic M, Lachmann B (1990) Haemodynamic and neurohumoral effects of xenon anaesthesia. A comparison with nitrous oxide. *Anaesthesia* 45(4):273–278
14. Luttrupp HH, Romner B, Perhag L, Eskilsson J, Fredriksen S, Werner O (1993) Left ventricular performance and cerebral haemodynamics during xenon anaesthesia. A transoesophageal echocardiography and transcranial Doppler sonography study. *Anaesthesia* 48(12):1045–1049
15. Rossaint R, Reyle-Hahn M, Schulte Am Esch J, Scholz J, Scherpereel P, Vallet B, Giunta F, Del Turco M et al (2003) Multicenter randomized comparison of the efficacy and safety of xenon and isoflurane in patients undergoing elective surgery. *Anesthesiology* 98(1):6–13
16. Coburn M, Kunitz O, Baumert JH, Hecker K, Haaf S, Zühlsdorff A, Becker T, Rossaint R (2005) Randomized controlled trial of the haemodynamic and recovery effects of xenon or propofol anaesthesia. *Br J Anaesth* 94(2):198–202
17. Wappler F, Rossaint R, Baumert J, Scholz J, Tonner PH, van Aken H, Berendes E, Klein J et al (2007) Multicenter randomized comparison of xenon and isoflurane on left ventricular function in patients undergoing elective surgery. *Anesthesiology* 106(3):463–471
18. Dingley J, King R, Hughes L, Terblanche C, Mahon S, Hepp M, Youhana A, Watkins A (2001) Exploration of xenon as a potential cardiostable sedative: a comparison with propofol after cardiac surgery. *Anaesthesia* 56(9):829–835

- 503 19. Goto T, Hanne P, Ishiguro Y, Ichinose F, Niimi Y, Morita S (2004) Cardiovascular effects of xenon and nitrous oxide in patients during
504 fentanyl-midazolam anaesthesia. *Anaesthesia* 59(12):1178–1183 552
- 505 20. Lockwood GG, Franks NP, Downie NA, Taylor KM, Maze M (2006) Feasibility and safety of delivering xenon to patients under-
506 going coronary artery bypass graft surgery while on cardiopulmo-
507 nary bypass: phase I study. *Anesthesiology* 104(3):458–465 553
- 508 21. Baumert JH, Hecker KE, Hein M, Reyle-Hahn M, Horn NA, Rossaint R (2005) Effects of xenon anaesthesia on the circulatory
509 response to hypoventilation. *Br J Anaesth* 95(2):166–171 554
- 510 22. Rex S, Schaefer W, Meyer PH, Rossaint R, Boy C, Setani K, Büll U, Baumert JH (2006) Positron emission tomography study of re-
511 gional cerebral metabolism during general anesthesia with xenon in
512 humans. *Anesthesiology* 105(5):936–943 555
- 513 23. Laitio RM, Kaisti KK, Långsjö JW, Aalto S, Salmi E, Maksimow A, Aantaa R, Oikonen V et al (2007) Effects of xenon anesthesia on
514 cerebral blood flow in humans: a positron emission tomography
515 study. *Anesthesiology* 106(6):1128–1133 556
- 516 24. Laitio RM, Långsjö JW, Aalto S, Kaisti KK, Salmi E, Maksimow A, Aantaa R, Oikonen V et al (2009) The effects of xenon anesthe-
517 sia on the relationship between cerebral glucose metabolism and
518 blood flow in healthy subjects: a positron emission tomography
519 study. *Anesth Analg* 108(2):593–600 557
- 520 25. Giller CA, Purdy P, Lindstrom WW (1990) Effects of inhaled stable
521 xenon on cerebral blood flow velocity. *AJNR Am J Neuroradiol* 11(1):177–182 558
- 522 26. Azzopardi D, Robertson NJ, Kapetanakis A, Griffiths J, Rennie JM, Mathieson SR, Edwards AD (2013) Anticonvulsant effect of xenon
523 on neonatal asphyxial seizures. *Arch Dis Child Fetal Neonatal Ed* 98(5):F437–F439 559
- 524 27. Choi DW, Koh JY, Peters S (1988) Pharmacology of glutamate
525 neurotoxicity in cortical cell culture: attenuation by NMDA antag-
526 onists. *J Neurosci* 8(1):185–196 560
- 527 28. Wilhelm S, Ma D, Maze M, Franks NP (2002) Effects of xenon on
528 in vitro and in vivo models of neuronal injury. *Anesthesiology* 96(6):1485–1491 561
- 529 29. Petzelt C, Blom P, Schmehl W, Müller J, Kox WJ (2003) Prevention
530 of neurotoxicity in hypoxic cortical neurons by the noble gas xenon.
531 *Life Sci* 72(17):1909–1918 562
- 532 30. David HN, Leveille F, Chazalviel L, MacKenzie ET, Buisson A, Lemaire M, Abraini JH (2003) Reduction of ischemic brain damage
533 by nitrous oxide and xenon. *J Cereb Blood Flow Metab* 23(10):
534 1168–1173 563
- 535 31. Nagata A, Nakao Si S, Nishizawa N, Masuzawa M, Inada T, Murao K, Miyamoto E, Shingu K (2001) Xenon inhibits but N(2)O en-
536 hances ketamine-induced c-Fos expression in the rat posterior cin-
537 gulate and retrosplenial cortices. *Anesth Analg* 92(2):362–368 564
- 538 32. Ma D, Lim T, Xu J, Tang H, Wan Y, Zhao H, Hossain M, Maxwell PH et al (2009) Xenon preconditioning protects against renal
539 ischemic-reperfusion injury via HIF-1 α activation. *J Am Soc Nephrol* 20(4):713–720 565
- 540 33. Stoppe C, Ney J, Brenke M, Goetzenich A, Emontzpohl C, Schälte G, Grottko O, Moeller M et al (2016) Sub-anesthetic xenon in-
541 creases erythropoietin levels in humans: a randomized controlled
542 trial. *Sports Med* 46(11):1753–1766 566
- 543 34. Albers GW, Goldstein LB, Hess DC, Wechsler LR, Furie KL, Gorelick PB, Hurn P, Liebeskind DS et al (2011) Stroke
544 Treatment Academic Industry Roundtable (STAIR) recommenda-
545 tions for maximizing the use of intravenous thrombolytics and
546 expanding treatment options with intra-arterial and neuroprotective
547 therapies. *Stroke* 42(9):2645–2650 567
- 548 35. Fries M, Nolte KW, Coburn M, Rex S, Timper A, Kottmann K, Siepmann K, Häusler M et al (2008) Xenon reduces
549 neurohistopathological damage and improves the early neurologi-
550 cal deficit after cardiac arrest in pigs. *Crit Care Med* 36(8):2420–
551 2426 568
- 552 36. Fries M, Brücken A, Çizen A, Westerkamp M, Löwer C, Deike-Glindemann J, Schnorrenberger NK, Rex S et al (2012) Combining
553 xenon and mild therapeutic hypothermia preserves neurological
554 function after prolonged cardiac arrest in pigs. *Crit Care Med* 40(4):1297–1303 569
- 555 37. Ma D, Hossain M, Chow A, Arshad M, Battson RM, Sanders RD, Mehmet H, Edwards AD et al (2005) Xenon and hypothermia com-
556 bine to provide neuroprotection from neonatal asphyxia. *Ann Neurol* 58(2):182–193 570
- 557 38. Martin JL, Ma D, Hossain M, Xu J, Sanders RD, Franks NP, Maze M (2007) Asynchronous administration of xenon and hypothermia
558 significantly reduces brain infarction in the neonatal rat. *Br J Anaesth* 98(2):236–240 571
- 559 39. Arola OJ, Laitio RM, Roine RO, Grönlund J, Saraste A, Pietilä M, Airaksinen J, Perttilä J et al (2013) Feasibility and cardiac safety of
560 inhaled xenon in combination with therapeutic hypothermia follow-
561 ing out-of-hospital cardiac arrest. *Crit Care Med* 41(9):2116–2124 572
- 562 40. Laitio R, Hynninen M, Arola O, Virtanen S, Parkkola R, Saunavaara J, Roine RO, Grönlund J et al (2016) Effect of inhaled
563 xenon on cerebral white matter damage in comatose survivors of
564 out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA* 315(11):1120–1128 573
- 565 41. Arola O, Saraste A, Laitio R, Airaksinen J, Hynninen M, Bäcklund M, Ylikoski E, Wennervirta J et al (2017) Inhaled xenon attenuates
566 myocardial damage in comatose survivors of out-of-hospital cardiac
567 arrest: the Xe-Hypotheca trial. *J Am Coll Cardiol* 70(21):2652–
568 2660 574

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES.

- Q1. Please check if the corresponding author's telecommunications data are correctly captured/ indicated.
- Q2. Please check if the affiliations are presented correctly.
- Q3. An abstract section is desired for this manuscript, but none is provided in the source files. Please then consider providing an appropriate abstract that substantively meets the requirements declared in the "Instructions for Authors" page of this journal.
- Q4. Keywords are required. The following are suggested: xenon, neuroprotection, medical application. Please check if appropriate; otherwise, please provide.
- Q5. Please check if the section headings are assigned to appropriate levels.
- Q6. The sentence "Following the discovery that xenon inhibits..." has been modified. Please check if the intended meaning was retained.
- Q7. A citation for Fig. 1 was inserted in the sentence "The Kaplan–Meier survival estimate (panel A)..." Please check if appropriate. Otherwise, please advise.